




Contents lists available at ScienceDirect

Advances in Applied Mathematics

www.elsevier.com/locate/yaama

Characterizing weak compatibility in terms of weighted quartets

Stefan Grünewald^{a,b}, Katharina T. Huber^c, Vincent Moulton^c,
Charles Semple^d, Andreas Spillner^{c,*}

^a CAS-MPG Partner Institute for Computational Biology, Chinese Academy of Sciences, Shanghai 200031, China

^b Max-Planck-Institute for Mathematics in the Sciences, 04103 Leipzig, Germany

^c School of Computing Sciences, University of East Anglia, Norwich, NR4 7TJ, UK

^d Department of Mathematics and Statistics, University of Canterbury, Christchurch, New Zealand

ARTICLE INFO

Article history:

Received 8 May 2008

Accepted 5 July 2008

Available online 14 November 2008

MSC:

92D15

Keywords:

Phylogenetics

Quartet

Weak compatibility

ABSTRACT

In phylogenetics there are various methods available for understanding the evolutionary history of a set of species based on the analysis of its 4-element subsets. Guided by biological data, such techniques usually require the initial computation of a quartet-weight function, i.e., a function that assigns a weight to each bipartition of each 4-element subset into two parts of size two, from which a phylogenetic tree or network is subsequently deduced. It is therefore of interest to characterize quartet-weight functions that correspond precisely to phylogenetic trees or networks. Recently, such characterizations have been presented for phylogenetic trees. Here we provide a 5-point condition for characterizing more general structures called weakly compatible split systems. Such split systems underly the construction of split networks, a special class of phylogenetic networks. This 5-point condition also yields a new characterization of quartet-weight functions that correspond to phylogenetic trees.

© 2008 Elsevier Inc. All rights reserved.

* Corresponding author.

E-mail addresses: stefan@picb.ac.cn (S. Grünewald), katharina.huber@cmp.uea.ac.uk (K.T. Huber), vincent.moulton@cmp.uea.ac.uk (V. Moulton), c.semple@math.canterbury.ac.nz (C. Semple), aspillner@cmp.uea.ac.uk (A. Spillner).

1. Introduction

Reconstructing evolutionary trees and, more generally, phylogenetic networks, is an important problem in evolutionary biology (see e.g. [9,12,17]). Formally speaking, for a set X of species, an evolutionary or *phylogenetic* (X)-tree T is a (graph theoretical) tree with leaf set X , no degree 2 vertices, and a weight function that assigns a non-negative weight to each edge of T . An example of such a tree is given in Fig. 1(a). The theory of such trees is well developed [18], and several methods are available for reconstructing them from biological data [12,17].

Any phylogenetic tree T may be encoded in terms of the subtrees T' of T that are spanned by the 4-element subsets of X [18, p. 130], cf. Fig. 1(b), and several methods for tree reconstruction rely on this fact (see e.g. [13,19,22]). With this in mind, let $\mathcal{Q}(X)$ denote the set of all bipartitions of the form $a_1a_2|b_1b_2$, where a_1, a_2, b_1, b_2 are distinct elements of X , i.e., $\mathcal{Q}(X)$ is the set of *quartets* on X . Then, for every quartet $a_1a_2|b_1b_2$, T induces weight $u(a_1a_2|b_1b_2)$ corresponding to the total weight of those edges in the subtree T' of T spanned by $\{a_1, a_2, b_1, b_2\}$ that are neither on the path from a_1 to a_2 nor on the path from b_1 to b_2 (see e.g. Fig. 1(b)). In particular, we obtain a *quartet-weight function*, i.e. a map $u : \mathcal{Q}(X) \rightarrow \mathbb{R}_{\geq 0}$.

As we have seen, it is straightforward to associate a quartet-weight function to a phylogenetic tree, but it is less obvious precisely which quartet-weight functions arise in this way. Even so, Dress and Erdős recently characterized those quartet-weight functions associated to *binary* phylogenetic trees [11] (that is, phylogenetic trees in which every internal vertex has degree 3) and Grünwald et al. [14] subsequently presented a characterization for phylogenetic trees in general (see also [1] and [7,8] for related results in the context of unweighted trees). In this paper we are interested in characterizing quartet-weight functions associated to structures that generalize phylogenetic trees.

To present our main result we first recall some additional facts concerning phylogenetic trees. To any edge e in a phylogenetic X -tree T we can associate a bipartition or *split* of X (see e.g. Fig. 1(c)). In particular, we obtain a *split-weight function*, i.e. a map w from the set $\Sigma(X)$ of all splits of X to $\mathbb{R}_{\geq 0}$, that assigns to each split of X associated to edge e of T the weight of e , and to all other splits weight 0. A fundamental result in phylogenetics [6] implies that phylogenetic trees correspond to split-weight functions w whose support, $\text{supp}(w) = \{S \in \Sigma(X) : w(S) > 0\}$, is *compatible* (i.e., for any two splits $A_1|B_1, A_2|B_2$ in $\text{supp}(w)$ at least one of the intersections $A_1 \cap A_2, A_1 \cap B_2, B_1 \cap A_2, B_1 \cap B_2$ is empty). Therefore, since any split-weight function w induces a quartet-weight function u_w defined by

$$u_w(a_1a_2|b_1b_2) = \sum_{\substack{A|B \in \Sigma(X), \\ \{a_1, a_2\} \subseteq A, \{b_1, b_2\} \subseteq B \text{ or } \{a_1, a_2\} \subseteq B, \{b_1, b_2\} \subseteq A}} w(A|B), \quad (1)$$

the above mentioned results in [11,14] can be regarded as characterizations of quartet-weight functions u for which there exists a split-weight function w with $u = u_w$ such that $\text{supp}(w)$ is compatible.

Here, we shall characterize quartet-weight functions u for which there exists a split-weight function w with $u = u_w$ such that $\text{supp}(w)$ is *weakly compatible* (i.e., for any three splits $A_1|B_1, A_2|B_2, A_3|B_3$ in $\text{supp}(w)$ at least one of the intersections $A_1 \cap A_2 \cap A_3, A_1 \cap B_2 \cap B_3, B_1 \cap A_2 \cap B_3, B_1 \cap B_2 \cap A_3$ is empty [2]). The concept of weak compatibility forms the basis for the construction of so-called *split*




Fig. 1. (a) A phylogenetic X -tree with $X = \{a_1, a_2, a_3, b_1, b_2, b_3, b_4\}$. (b) The subtree spanned by $\{a_1, a_2, b_1, b_2\}$. The induced weight of the quartet $a_1a_2|b_1b_2$ is 5, the total weight of the bold edges. (c) In this phylogenetic tree the split $a_1a_2|a_3a_4a_5$ is associated with edge e .

networks [3,10,15], a special class of labeled, weighted, graphs used to understand complex patterns of evolution [16] that generalize phylogenetic trees. Our main result is the following.

Theorem 1. Suppose that X is a finite set, $u : \mathcal{Q}(X) \rightarrow \mathbb{R}_{\geq 0}$ is a quartet-weight function, and, for $q \in \{\leq 1, = 1, \leq 2, = 2\}$, consider the following properties:

- (W1)^q For every 4 distinct elements $a, b, c, d \in X$ at most 1 (precisely 1, at most 2, precisely 2) of the quantities $u(ab|cd)$, $u(ac|bd)$ and $u(ad|bc)$ are non-zero.
 (W2) For every 5 distinct elements a_1, a_2, b_1, b_2, x in X ,

$$u(a_1 a_2 | b_1 b_2) = \min \left\{ \begin{array}{l} u(a_1 a_2 | b_1 b_2) \\ u(a_1 x | b_1 b_2) \\ u(a_2 x | b_1 b_2) \end{array} \right\} + \min \left\{ \begin{array}{l} u(a_1 a_2 | b_1 b_2) \\ u(a_1 a_2 | b_1 x) \\ u(a_1 a_2 | b_2 x) \end{array} \right\}.$$

Then the following statements hold.

- (A) There exists a split-weight function w with $u = u_w$ and $\text{supp}(w)$ weakly compatible if and only if u satisfies (W1) ^{≤ 2} and (W2).
 (B) There exists a split-weight function w with $u = u_w$ and $\text{supp}(w)$ compatible if and only if u satisfies (W1) ^{≤ 1} and (W2).
 (C) There exists a split-weight function w with $u = u_w$ and $\text{supp}(w)$ maximal (and, therefore, maximum) compatible if and only if u satisfies (W1) ^{$= 1$} and (W2).

Note that (B) and (C) are alternative characterizations to those given in [14] and [11] for when a quartet-weight function arises from a phylogenetic tree and a binary phylogenetic tree, respectively. Furthermore, (A) can be viewed as a generalization of Bandelt and Dress' 6-point condition in [4] that essentially characterizes quartet sets of the form $\text{supp}(u_w) = \{q \in \mathcal{Q}(X) : u_w(q) > 0\}$, w a split-weight function with the property that $\text{supp}(w)$ is weakly compatible. Note that, in contrast to (A), the induced weights of the quartets in $\text{supp}(u_w)$ are ignored in [4] and, therefore, also the precise weights of the splits in $\text{supp}(w)$ are not important. This results in a loss of information that is illustrated by an example given in [4, p. 126] which shows that no characterization of these quartet sets is possible in terms of an i -point condition with $i \leq 5$.

Note also that if a quartet-weight function u satisfies (W2) and (W1) ^{$= 2$} , then one can show – using a completely analogous argument as in the proof of characterization (C) given below – that there exists a split-weight function w with $u = u_w$ and $\text{supp}(w)$ maximal weakly compatible (although this does not necessarily imply that $\text{supp}(w)$ is maximum weakly compatible [2, p. 70]). However, the converse statement does not hold. For example, if $X = \{a, b, c, d, e, f\}$, and w is the split-weight function on $\Sigma(X)$ that assigns weight 1 to each of the following splits of X : $ab|cdef$, $abe|cdf$, $abef|cd$, $ad|bcef$, $adf|bce$, $adef|bc$ and $x|X - x$ for every $x \in X$, and 0 to every other split, then it can be easily checked that $\text{supp}(w)$ is maximal weakly compatible, although for the 4-element subset $\{b, d, e, f\}$ only $u_w(be|df)$ is non-zero.

The rest of the paper is organized as follows. In Section 2, we introduce some basic notation. In Section 3, we prove some useful results concerning quartet-weight functions, and use these to prove that characterization (A) holds. In Section 4, we prove that characterizations (B) and (C) hold. We conclude in Section 5 with some observations concerning the characterization of quartet-weight functions which correspond to split-weight functions whose support is *circular*, a property that generalizes compatibility but that is more restrictive than weak compatibility [2]. In particular, we show that it is not possible to characterize such quartet-weight functions by any i -point condition, $i \in \mathbb{N}$.

2. Preliminaries

For any two non-empty subsets A and B of X with the property that $A \cap B = \emptyset$, we call $A|B$ a *partial split* of X . In particular, a quartet is a partial split. We denote the set of all partial splits $A|B$ of X with $\min\{|A|, |B|\} \geq 2$ by $\Sigma_p^*(X)$. For any two partial splits $A_1|B_1$ and $A_2|B_2$ of X , we say that

$A_2|B_2$ extends $A_1|B_1$, denoted by $A_2|B_2 \succ A_1|B_1$, if $A_2 \supseteq A_1$ and $B_2 \supseteq B_1$, or $A_2 \supseteq B_1$ and $B_2 \supseteq A_1$. For $A \subseteq X$ and $x \in X - A$, we use $A + x$ to denote $A \cup \{x\}$.

Now let $\mathfrak{U}(X)$ denote the set of quartet-weight functions on $\mathcal{Q}(X)$ and $\mathfrak{W}(X)$ the set of split-weight functions on $\Sigma(X)$. Recall that a split $A|B$ of X is called *trivial* if $\min\{|A|, |B|\} = 1$. Note that for every $\mathfrak{w} \in \mathfrak{W}(X)$ only the non-trivial splits, i.e., the splits in $\Sigma^*(X) = \{A|B \in \Sigma(X) : \min\{|A|, |B|\} \geq 2\}$, contribute to $u_{\mathfrak{w}}$ in Eq. (1).

Note that every $\mathfrak{w} \in \mathfrak{W}(X)$ induces a *distance function* $D_{\mathfrak{w}}$ as follows:

$$D_{\mathfrak{w}}(x, y) := \sum_{S \in \Sigma(X), S \succ x|y} \mathfrak{w}(S)$$

for every $(x, y) \in X \times X$, i.e., a symmetric map $D_{\mathfrak{w}} : X \times X \rightarrow \mathbb{R}_{\geq 0}$ with the property that $D(x, x) = 0$ for every $x \in X$. This function is always a (*pseudo*-)metric, that is, it satisfies the triangle inequality $D_{\mathfrak{w}}(x, z) \leq D_{\mathfrak{w}}(x, y) + D_{\mathfrak{w}}(y, z)$ for all $x, y, z \in X$. *Split decomposition* [2] reverses this process. In particular, given a distance function D , a weight function $\alpha = \alpha_D$ on the set of all partial splits of X is defined as follows:

$$\alpha(A|B) := \frac{1}{2} \min_{\substack{a_1, a_2 \in A \\ b_1, b_2 \in B}} \left(\max \left\{ \begin{array}{l} D(a_1, b_1) + D(a_2, b_2), \\ D(a_1, b_2) + D(a_2, b_1), \\ D(a_1, a_2) + D(b_1, b_2) \end{array} \right\} - D(a_1, a_2) - D(b_1, b_2) \right)$$

for every partial split $A|B$ of X . Obviously, this yields a split-weight function \mathfrak{w}_D by restricting α to $\Sigma(X)$.

Central to the theory of split decomposition are the so-called *totally split-decomposable metrics*. Such a metric D on X can be written as $D = D_{\mathfrak{w}}$ where $\mathfrak{w} \in \mathfrak{W}(X)$ has the property that $\text{supp}(\mathfrak{w})$ is weakly compatible. For brevity, we will call $\mathfrak{w} \in \mathfrak{W}(X)$ *weakly compatible* if $\text{supp}(\mathfrak{w})$ is weakly compatible. Note that for a totally split-decomposable metric D there exists a unique weakly compatible split-weight function \mathfrak{w} with the property that $D = D_{\mathfrak{w}}$ and, in addition, for every split $S \in \Sigma(X)$ we have $\alpha(S) = \mathfrak{w}(S)$ [2, Theorem 3].

Finally, given a quartet-weight function $u \in \mathfrak{U}(X)$, we define a weight function γ_u on the set of all partial splits of X by

$$\gamma_u(A|B) := \min\{u(q) : q \in \mathcal{Q}(X), A|B \succ q\}$$

where $A|B \in \Sigma_p^*(X)$, and $\gamma_u(A|B) = 0$ for all other partial splits of X . In case the quartet-weight function u is understood from the context, we will write $\gamma(A|B)$ rather than $\gamma_u(A|B)$. The restriction of γ_u to $\Sigma(X)$ is denoted by \mathfrak{w}_u . Note that property (W2) can now be written more concisely as

$$\gamma_u(a_1 a_2 | b_1 b_2) = \gamma_u(a_1 a_2 x | b_1 b_2) + \gamma_u(a_1 a_2 | b_1 b_2 x)$$

for every five distinct elements a_1, a_2, b_1, b_2, x in X . We conclude by rephrasing a simple but useful fact from [2, p. 60].

Fact 2. Let $\mathfrak{w} \in \mathfrak{W}(X)$. Then \mathfrak{w} is weakly compatible if and only if $u_{\mathfrak{w}}$ satisfies (W1) $^{\leq 2}$.

3. Proof of characterization (A)

The proof is organized as follows. We first show that quartet-weight functions that are induced by a weakly compatible split-weight function always satisfy (W1) $^{\leq 2}$ and (W2) (Lemma 3). The converse could be shown by proving analogous results on split decomposition theory appearing in [2] for quartet-weight functions. However, we will use a more direct approach: we first show that it suffices to prove a key equality (Lemma 4(ii)) and then establish that equality in Lemma 5.

Lemma 3. If $u \in \mathcal{U}(X)$ can be written as $u = u_{\mathfrak{w}}$ for some weakly compatible $\mathfrak{w} \in \mathcal{W}(X)$, then u satisfies properties (W1) $^{\leq 2}$ and (W2).

Proof. Let $\mathfrak{w} \in \mathcal{W}(X)$ be weakly compatible. Then, by Fact 2, $u = u_{\mathfrak{w}}$ satisfies (W1) $^{\leq 2}$. To show that u satisfies also (W2), put $\alpha = \alpha_{D_{\mathfrak{w}}}$ and $\gamma = \gamma_{u_{\mathfrak{w}}}$. As a first step, we show that $\alpha(A|B) = \gamma(A|B)$ for every partial split $A|B \in \Sigma_p^*(X)$.

To this end, consider an arbitrary partial split $A|B \in \Sigma_p^*(X)$. If $\alpha(A|B) > 0$, then, since $D_{\mathfrak{w}}$ is totally split decomposable, by [2, Theorem 6(ii)] we have $\alpha(A|B) = \sum_{S \in \Sigma(X), S \succ A|B} \mathfrak{w}(S)$. If $\alpha(A|B) = 0$, then it follows from the definition of α that $\mathfrak{w}(S) = 0$ for every split S of X such that $S \succ A|B$. Hence, $\alpha(q) = \sum_{S \in \Sigma(X), S \succ q} \mathfrak{w}(S) = u_{\mathfrak{w}}(q)$ for every $q \in \mathcal{Q}(X)$. Moreover, since $D_{\mathfrak{w}}$ is a metric, it follows from an observation in [2, p. 54] that $\alpha(A|B) = \min\{\alpha(q) : q \in \mathcal{Q}(X), A|B \succ q\}$, which, by the above, equals $\min\{u_{\mathfrak{w}}(q) : q \in \mathcal{Q}(X), A|B \succ q\} = \gamma(A|B)$ for every partial split $A|B$ in $\Sigma_p^*(X)$.

We now show that $u_{\mathfrak{w}}$ satisfies property (W2). Since $\alpha(A|B) = \gamma(A|B)$ for every partial split $A|B \in \Sigma_p^*(X)$, this follows immediately from [2, Theorem 6(iii)] which states that $\alpha(a_1 a_2 | b_1 b_2) = \alpha(a_1 a_2 x | b_1 b_2) + \alpha(a_1 a_2 | b_1 b_2 x)$ for any 5 distinct elements $a_1, a_2, b_1, b_2, x \in X$. \square

The next lemma establishes that to show that the converse of Lemma 3 holds, it suffices to show that Eq. (3) below holds.

Lemma 4. Let $u \in \mathcal{U}(X)$ satisfy properties (W1) $^{\leq 2}$ and (W2).

(i) For every partial split $A|B \in \Sigma_p^*(X)$ and every $x \in X - (A \cup B)$,

$$\gamma(A|B) \geq \gamma(A + x|B) + \gamma(A|B + x). \quad (2)$$

(ii) If

$$\gamma(A|B) = \gamma(A + x|B) + \gamma(A|B + x) \quad (3)$$

for every partial split $A|B \in \Sigma_p^*(X)$ and every $x \in X - (A \cup B)$, then $u = u_{\mathfrak{w}}$ for some weakly compatible $\mathfrak{w} \in \mathcal{W}(X)$.

Proof. (i) Let $A|B \in \Sigma_p^*(X)$ and $x \in X - (A \cup B)$. Choose two distinct elements $a_1, a_2 \in A$ and two distinct elements $b_1, b_2 \in B$ such that $\gamma(A|B) = u(a_1 a_2 | b_1 b_2)$ holds. Then

$$\begin{aligned} \gamma(A + x|B) + \gamma(A|B + x) &\leq \gamma(a_1 a_2 x | b_1 b_2) + \gamma(a_1 a_2 | b_1 b_2 x) \\ &= u(a_1 a_2 | b_1 b_2) = \gamma(A, B), \end{aligned}$$

where the second-to-last equality follows from property (W2).

(ii) First recall that the split-weight function $\mathfrak{w} = \mathfrak{w}_u$ is defined as the restriction of γ to $\Sigma(X)$. Since u satisfies property (W1) $^{\leq 2}$, it follows by Fact 2 that \mathfrak{w} is weakly compatible. Thus, it suffices to show that $u = u_{\mathfrak{w}}$. To do this, we use induction on the size k of $X - (A \cup B)$, and the induction hypothesis that

$$\gamma(A|B) = \sum_{S \in \Sigma(X), S \succ A|B} \mathfrak{w}(S)$$

holds for every partial split $A|B \in \Sigma_p^*(X)$.

The base case $k = 0$ states that $\gamma(S) = \mathfrak{w}(S)$ for every $S \in \Sigma(X)$. But this holds by definition.

Now suppose $k > 0$ and suppose $A|B \in \Sigma_p^*(X)$. Then there exists some $x \in X - (A \cup B)$. Using Eq. (3) it follows by induction that

$$\begin{aligned}
\gamma(A|B) &= \gamma(A+x|B) + \gamma(A|B+x) \\
&= \sum_{S \in \Sigma(X), S \succ A+x|B} \mathfrak{w}(S) + \sum_{S \in \Sigma(X), S \succ A|B+x} \mathfrak{w}(S) \\
&= \sum_{S \in \Sigma(X), S \succ A|B} \mathfrak{w}(S),
\end{aligned}$$

and so $u(q) = \gamma(q) = \sum_{S \in \Sigma(X), S \succ q} \mathfrak{w}(S)$ for every quartet $q \in \mathcal{Q}(X)$, as required. \square

The remainder of this section is devoted to the proof of the following lemma which establishes that properties (W1) $^{\leq 2}$ and (W2) imply Eq. (3).

Lemma 5. *Let $u \in \mathcal{U}(X)$ satisfy properties (W1) $^{\leq 2}$ and (W2). Then Eq. (3) holds for every partial split $A|B \in \Sigma_p^*(X)$ and every $x \in X - (A \cup B)$.*

To prove this lemma we use induction on $k := |A \cup B|$. Note that the base case $k = 4$ of the induction follows directly from property (W2). The remainder of the inductive proof is divided into two parts. In Part 1 we show that Eq. (3) holds for $k = 5$. This is the main part of the proof and is somewhat technical. In Part 2 we establish that Eq. (3) holds for $k \geq 6$. The following simple fact will be used several times in our proof.

Fact 6. Let $A|B \in \Sigma_p^*(X)$ and $x \in X - (A \cup B)$ be such that $\gamma(A|B) > \gamma(A+x|B)$. Then there exist $a \in A$ and $b_1, b_2 \in B$, $b_1 \neq b_2$, such that $\gamma(A+x|B) = u(ax|b_1b_2)$.

Part 1: $k = 5$. For the purpose of contradiction, we assume that there exists a partial split $A|B \in \Sigma_p^*(X)$, $|A| = 2$ and $|B| = 3$, and $x \in X - (A \cup B)$ such that

$$\gamma(A|B) > \gamma(A+x|B) + \gamma(A|B+x). \quad (4)$$

Note that (4) implies that $\gamma(A|B) > 0$ and, therefore, $u(q) > 0$ for every quartet q that is extended by $A|B$. Starting with the above assumption, we generate additional partial splits $A'|B'$, $|A'| = 2$ and $|B'| = 3$, satisfying inequality (4) until we obtain a contradiction to (W1) $^{\leq 2}$. We use the following lemma to generate these additional splits.

Lemma 7. *Suppose $A|B \in \Sigma_p^*(X)$, with $|A| = 2$ and $|B| = 3$, and $x \in X - (A \cup B)$ is such that inequality (4) holds. Then there exist precisely two elements $b \in B$ such that*

(i)

$$\begin{aligned}
\gamma(A+x|B-b) &> \gamma(A+x+b|B-b) + \gamma(A+x|B) \quad \text{and} \\
\gamma(A|B+x-b) &= \gamma(A|B+x),
\end{aligned}$$

and there exists precisely one element $b \in B$ such that

(ii)

$$\begin{aligned}
\gamma(A+x|B-b) &= \gamma(A+x|B) \quad \text{and} \\
\gamma(A|B+x-b) &> \gamma(A+b|B+x-b) + \gamma(A|B+x).
\end{aligned}$$

Moreover, no element in B satisfies both (i) and (ii).

Proof. First note that since $\gamma(A|B) > \gamma(A|B+x)$, by Fact 6 there exist at least two elements $b \in B$ such that $\gamma(A|B+x-b) = \gamma(A|B+x)$. Also since $\gamma(A|B) > \gamma(A+x|B)$, again by Fact 6, there exists at least one element $b \in B$ such that $\gamma(A+x|B-b) = \gamma(A+x|B)$. Clearly, there is no $b \in B$ such that $\gamma(A|B+x-b) = \gamma(A|B+x)$ and $\gamma(A+x|B-b) = \gamma(A+x|B)$ since otherwise, applying the induction hypothesis to $A|B-b$, we have

$$\begin{aligned}\gamma(A|B) &\leq \gamma(A|B-b) = \gamma(A+x|B-b) + \gamma(A|B+x-b) \\ &= \gamma(A+x|B) + \gamma(A|B+x)\end{aligned}$$

contradicting (4). Next note that there is no $b \in B$ such that

$$\begin{aligned}\gamma(A+x|B-b) &= \gamma(A+x+b|B-b) + \gamma(A+x|B) \quad \text{and} \\ \gamma(A|B+x-b) &= \gamma(A|B+x).\end{aligned}$$

To see this, suppose it were otherwise and note that again by applying the induction hypothesis to $A|B-b$ we have

$$\begin{aligned}\gamma(A|B-b) &= \gamma(A+x|B-b) + \gamma(A|B+x-b) \quad \text{as well as} \\ \gamma(A|B-b) &= \gamma(A+b|B-b) + \gamma(A|B).\end{aligned}$$

But then

$$\gamma(A+b|B-b) + \gamma(A|B) = \gamma(A+x+b|B-b) + \gamma(A+x|B) + \gamma(A|B+x)$$

which implies $\gamma(A|B) \leq \gamma(A+x|B) + \gamma(A|B+x)$ since $\gamma(A+x+b|B-b) \leq \gamma(A+b|B-b)$. But this contradicts (4). Similarly we can show that there is no $b \in B$ such that

$$\begin{aligned}\gamma(A+x|B-b) &= \gamma(A+x|B) \quad \text{and} \\ \gamma(A|B+x-b) &= \gamma(A+b|B+x-b) + \gamma(A|B+x).\end{aligned}$$

This, together with Lemma 4(i), completes the proof of the lemma. \square

We now apply Lemma 7 for the generation of additional partial splits $A'|B'$ with $\gamma(A'|B') > 0$. Let $A = \{a_1, a_2\}$ and $B = \{b_1, b_2, b_3\}$. Recall that we assume $\gamma(a_1a_2|b_1b_2b_3) > \gamma(a_1a_2x|b_1b_2b_3) + \gamma(a_1a_2|b_1b_2b_3x)$. Applying Lemma 7, we can assume by symmetry and without loss of generality that

$$\begin{aligned}\gamma(a_1a_2x|b_1b_2) &> \gamma(a_1a_2b_3x|b_1b_2) + \gamma(a_1a_2x|b_1b_2b_3), \\ \gamma(a_1a_2x|b_2b_3) &> \gamma(a_1a_2b_1x|b_2b_3) + \gamma(a_1a_2x|b_1b_2b_3) \quad \text{and} \\ \gamma(a_1a_2|b_1b_3x) &> \gamma(a_1a_2b_2|b_1b_3x) + \gamma(a_1a_2|b_1b_2b_3x).\end{aligned}$$

(Note that this also determines uniquely the remaining equalities that must hold by Lemma 7.) Similarly, applying Lemma 7 to the partial split $b_1b_2|a_1a_2x$, we can again assume by symmetry and without loss of generality that

$$\gamma(b_1b_2b_3|a_1x) > \gamma(a_2b_1b_2b_3|a_1x) + \gamma(b_1b_2b_3|a_1a_2x).$$

Now, by Lemma 7(ii), either

$$\gamma(b_1b_2|a_2b_3x) > \gamma(a_1b_1b_2|a_2b_3x) + \gamma(b_1b_2|a_1a_2b_3x)$$

or

$$\gamma(b_1b_2|a_1a_2b_3) > \gamma(b_1b_2x|a_1a_2b_3) + \gamma(b_1b_2|a_1a_2b_3x).$$

But $\gamma(b_1b_2b_3|a_1a_2) \neq \gamma(b_1b_2b_3|a_1a_2x)$ as $\gamma(a_1a_2|b_1b_2b_3) > \gamma(a_1a_2x|b_1b_2b_3) + \gamma(a_1a_2|b_1b_2b_3x)$, and so the first of these two inequalities must hold. Similarly, applying Lemma 7 to the partial split $b_2b_3|a_1a_2x$, implies

$$\gamma(b_2b_3|a_2b_1x) > \gamma(a_1b_2b_3|a_2b_1x) + \gamma(b_2b_3|a_1a_2b_1x),$$

and, applying Lemma 7 to the partial split $b_1b_2|a_2b_3x$ and then to the partial split $b_2b_3|a_2b_1x$, implies

$$\gamma(a_1b_1b_2|b_3x) > \gamma(a_1a_2b_1b_2|b_3x) + \gamma(a_1b_1b_2|a_2b_3x) \quad \text{and}$$

$$\gamma(a_1b_2b_3|b_1x) > \gamma(a_1a_2b_2b_3|b_1x) + \gamma(a_1b_2b_3|a_2b_1x).$$

Hence, since $\gamma(b_1b_2b_3|a_1x) > 0$, $\gamma(a_1b_1b_2|b_3x) > 0$ and $\gamma(a_1b_2b_3|b_1x) > 0$ and since $u(q) > 0$ for every quartet extended by $b_1b_2b_3|a_1x$, $a_1b_1b_2|b_3x$, and $a_1b_2b_3|b_1x$, we must have $u(a_1x|b_1b_3) > 0$, $u(a_1b_1|b_3x) > 0$ and $u(a_1b_3|b_1x) > 0$, contradicting (W1) $^{\leq 2}$. This completes the proof of Part 1 and so Eq. (3) holds for $k = 5$.

Part 2: $k \geq 6$. We first show that Eq. (3) holds for $k = 6$. Note that if $\gamma(A|B) = \gamma(A+x|B)$ or $\gamma(A|B) = \gamma(A|B+x)$, then $\gamma(A|B) = \gamma(A+x|B) + \gamma(A|B+x)$ by Lemma 4(i). So assume that $\gamma(A|B) > \gamma(A+x|B)$ and $\gamma(A|B) > \gamma(A|B+x)$, and consider the following two cases.

Case 1: $\max\{|A|, |B|\} = 4$. Without loss of generality assume that $|A| = 4$ and $|B| = 2$. By Fact 6, since $|A| = 4$, we can select $a \in A$ such that $\gamma(A+x-a|B) = \gamma(A+x|B)$ and $\gamma(A-a|B+x) = \gamma(A|B+x)$. Then

$$\begin{aligned} \gamma(A|B) &\leq \gamma(A-a|B) = \gamma(A+x-a|B) + \gamma(A-a|B+x) \\ &= \gamma(A+x|B) + \gamma(A|B+x) \end{aligned}$$

by (3) for $k = 5$. But then, by Lemma 4(i), $\gamma(A|B) = \gamma(A+x|B) + \gamma(A|B+x)$.

Case 2: $|A| = |B| = 3$. By Fact 6, since $|A| = 3$, we can select $a \in A$ such that $\gamma(A+x-a|B) = \gamma(A+x|B)$. By (3) for $k = 5$ and Case 1, we obtain

$$\begin{aligned} \gamma(A-a|B) &= \gamma(A+x-a|B) + \gamma(A-a|B+x) \\ &= \gamma(A+x-a|B) + \gamma(A|B+x) + \gamma(A-a|B+x+a), \end{aligned}$$

and, similarly,

$$\begin{aligned} \gamma(A-a|B) &= \gamma(A|B) + \gamma(A-a|B+a) \\ &= \gamma(A|B) + \gamma(A+x-a|B+a) + \gamma(A-a|B+x+a). \end{aligned}$$

It follows that

$$\gamma(A + x - a|B) + \gamma(A|B + x) = \gamma(A|B) + \gamma(A + x - a|B + a)$$

from which, by the choice of a ,

$$\gamma(A + x|B) + \gamma(A|B + x) \geq \gamma(A|B)$$

follows. But then, by Lemma 4(i), $\gamma(A|B) = \gamma(A + x|B) + \gamma(A|B + x)$. This completes the proof of (3) for $k = 6$.

So, suppose $k \geq 7$. But then $\max\{|A|, |B|\} \geq 4$, and so we can apply the same argument (using induction) as used in Case 1 for $k = 6$. This completes the proof of Part 2. \square

4. Proof of characterizations (B) and (C)

Proof of (B). Suppose $\mathfrak{w} \in \mathfrak{W}(X)$ with $\text{supp}(\mathfrak{w})$ compatible and $u = u_{\mathfrak{w}}$. Since every compatible split system is weakly compatible, it follows from characterization (A) that u satisfies $(W1)^{\leq 2}$ and $(W2)$. To see that u must satisfy even $(W1)^{\leq 1}$ assume for contradiction that there exist 4 distinct elements $a, b, c, d \in X$ such that at least two of the quantities $u(ab|cd)$, $u(ac|bd)$ and $u(ad|bc)$ are non-zero. Without loss of generality assume $u(ab|cd)$ and $u(ac|bd)$ are non-zero. But then, since quartets $ab|cd$ and $ac|bd$ must be extended by a split in $\text{supp}(\mathfrak{w})$, it follows that $\text{supp}(\mathfrak{w})$ is not compatible, a contradiction.

To prove the converse, assume that $u \in \mathfrak{U}(X)$ satisfies $(W1)^{\leq 1}$ and $(W2)$. Then u satisfies $(W1)^{\leq 2}$ and $(W2)$. Hence, by characterization (A), there exists a weakly compatible $\mathfrak{w} \in \mathfrak{W}(X)$ such that $u = u_{\mathfrak{w}}$. But now it follows directly from $(W1)^{\leq 1}$ that $\text{supp}(\mathfrak{w})$ must even be compatible. \square

Proof of (C). Suppose $\mathfrak{w} \in \mathfrak{W}(X)$ with $\text{supp}(\mathfrak{w})$ maximal compatible and $u = u_{\mathfrak{w}}$. By characterization (B) it remains to show that this implies $(W1)^{=1}$. But this is well known [7,8,11].

To see that the converse holds, suppose that $u \in \mathfrak{U}(X)$ satisfies $(W1)^{=1}$ and $(W2)$. By characterization (B), there exists $\mathfrak{w} \in \mathfrak{W}(X)$ with the property that $u = u_{\mathfrak{w}}$ and $\text{supp}(\mathfrak{w})$ is compatible. We may assume without loss of generality that $\text{supp}(\mathfrak{w})$ contains the trivial splits of X . Now assume for a contradiction that there exists a split $S' \in \Sigma^*(X) - \text{supp}(\mathfrak{w})$ such that $\text{supp}(\mathfrak{w}) + S'$ is still compatible, and define a split-weight function \mathfrak{w}' by $\mathfrak{w}'(S) = \mathfrak{w}(S)$ for every split $S \in \Sigma(X) - S'$ and $\mathfrak{w}'(S') = 1$. Since $\text{supp}(\mathfrak{w}')$ is compatible, by characterization (B), the quartet-weight function $u' = u_{\mathfrak{w}'}$ induced by \mathfrak{w}' must satisfy $(W1)^{\leq 1}$ and $(W2)$. Furthermore, since $\mathfrak{w}'(S') = \gamma_{u'}(S') > \gamma_u(S') = \mathfrak{w}(S') = 0$, there must exist a quartet $q \in \mathcal{Q}(X) - \text{supp}(u)$ such that q is extended by split S' . But since u satisfies $(W1)^{=1}$ and by construction $\text{supp}(u) \subseteq \text{supp}(u')$, this contradicts the fact that u' satisfies $(W1)^{\leq 1}$. \square

5. Circular split systems

We have seen how to characterize weakly compatible quartet-weight functions, functions that arise in the context of split networks [2,3]. An important subclass of these functions that are also widely used in this context are those corresponding to *circular split systems*. A split system $\Sigma' \subseteq \Sigma(X)$ is called *circular* if there exists an ordering x_1, x_2, \dots, x_n of X with the property that for every split $A|B \in \Sigma'$ there are $i, j \in \{1, \dots, n\}$, $i \leq j$, such that $A = \{x_i, \dots, x_j\}$ or $B = \{x_i, \dots, x_j\}$ [2]. Note that every compatible split system is circular, and that every maximum weakly compatible split system is (maximum) circular [2]. Circular split systems and the corresponding quartet-weight functions arise in the construction of *planar split networks* [5,13].

In view of our above results, it is natural to ask whether it is possible to give i -point characterizations for quartet-weight functions that are induced by split-weight functions whose support is circular. Note that Bandelt and Dress [2] characterized the quartet sets $\text{supp}(u_{\mathfrak{w}})$ that arise from a




Fig. 2. Examples of forbidden split systems. Elements in X are represented as dots and splits by curves or curve segments, for example, in (a), $x_5x_6 | x_1x_2x_3x_4$ and $x_1x_3x_5 | x_2x_4x_6$ are splits. The split system pictured in (a) is Ψ and the split system in (b) is Γ_9 .

split weight function \mathfrak{w} with the property that $\text{supp}(\mathfrak{w})$ is *maximum* circular by a 5-point condition (see also [21]). However, we shall now show that in general there is no such i -point characterization, $i \in \mathbb{N}$.

Given a split system $\Sigma' \subseteq \Sigma(X)$ and some subset $Y \subseteq X$, define the split system *induced by* Σ' on Y by $\Sigma'_Y = \{A \cap Y | B \cap Y : A | B \in \Sigma'\} \cap \Sigma(Y)$. In [20, p. 18], it is shown that a split system Σ cannot be circular if there is a 6-element subset $Y = \{x_1, x_2, \dots, x_6\} \subseteq X$ and $\Sigma' \subseteq \Sigma$ such that the split system induced by Σ' on Y is the split system Ψ in Fig. 2(a) or there is a k -element subset $Y = \{x_1, x_2, \dots, x_k\} \subseteq X$, $k \geq 4$, and $\Sigma' \subseteq \Sigma$ such that the split system induced by Σ' on Y is the split system

$$\Gamma_k = \left\{ \{x_i, x_{i+1}\} | X - \{x_i, x_{i+1}\} : 1 \leq i \leq k-2 \right\} \\ \cup \left\{ \{x_{k-1}, x_1\} | X - \{x_{k-1}, x_1\} \right\}$$

(see Fig. 2(b) where the split system Γ_9 is pictured). We will refer to the split systems Ψ and Γ_k , $k \geq 4$, as the *forbidden* split systems.

It follows immediately that no i -point condition, $i \in \mathbb{N}$, characterizes quartet-weight functions corresponding to split-weight functions with circular support. Even so, we next present a result of independent interest that implies that the above configurations are in some sense enough to characterize circular split systems.

Theorem 8. *A split system Σ on X is circular if and only if there are no subsets Σ' of Σ and Y of X such that the split system Σ'_Y is one of the forbidden split systems.*

Note that an alternative characterization of circular split systems that employs a set theoretical closure operation may be found in [20, Theorem 1.29]. The remainder of this section is devoted to the proof of Theorem 8. In view of the discussion above, it suffices to show that if Σ is *clean* on X , i.e. there are no subsets Σ' of Σ and Y of X such that the split system Σ'_Y is one of the forbidden split systems, then Σ is circular.

Assume for a contradiction that there exists a split system Σ on some set X such that Σ is clean on X but not circular. Fix such a Σ with $|X|$ minimal. Then it follows that $|X| \geq 4$, since every split system on a set with at most 3 elements is circular.

Now select an arbitrary element $z \in X$ and define $Z = X - \{z\}$. Note that the induced split system $\Sigma_{|Z}$ is clean on Z . Thus, since $n = |Z| < |X|$, by the minimality of $|X|$, there exists a circular ordering $\Theta = x_1, \dots, x_n$ of Z that is *compatible with* $\Sigma_{|Z}$, i.e., for every split $A | B \in \Sigma_{|Z}$ there are $i, j \in \{1, \dots, n\}$, $i \leq j$, such that $A = \{x_i, \dots, x_j\}$ or $B = \{x_i, \dots, x_j\}$. In the following, when dealing with indices taken from the set $\{1, 2, \dots, l\}$ for some integer $l \geq 1$, it will be convenient to allow also index $l+1$ and agree that the element indexed by $l+1$ is the same as the element indexed by 1.

Since the trivial splits of X are compatible with every ordering of X , we can assume without loss of generality that Σ does not contain any trivial splits. Then, for each split $S \in \Sigma$, we let A_S denote the element in S that does not contain z . Note that for every split $S \in \Sigma$ there exists some $S' \in \Sigma_{|Z}$ such that $A_S \in S'$. We continue the proof of Theorem 8 with the following lemma.

Lemma 9. *There are two splits S_1 and S_2 in Σ such that (shifting ordering Θ suitably if necessary)*

$$A_{S_1} = \{x_1, \dots, x_a\} \quad \text{and} \quad A_{S_2} = \{x_{b_1}, \dots, x_n, x_1, \dots, x_{b_2}\}$$

with $1 \leq b_2$, $b_2 + 2 \leq b_1$, $b_1 \leq a$, and $a < n$.

Proof. We divide our argument into two cases.

Case 1. There exists some $c \in \{1, \dots, n\}$ such that there is no split $S \in \Sigma$ with the property that $\{x_c, x_{c+1}\}$ is a subset of A_S . Then the ordering $\Theta' = x_1, \dots, x_c, z, x_{c+1}, \dots, x_n$ of X is compatible with Σ , contradicting our choice of Σ .

Case 2. For every $c \in \{1, \dots, n\}$ there exists a split $S \in \Sigma$ such that $\{x_c, x_{c+1}\}$ is a subset of A_S . Then there must exist splits S_1, \dots, S_l in Σ and elements z_1, \dots, z_l in Z , $l \geq 2$, such that for every $i \in \{1, \dots, l\}$ element z_i is contained in A_{S_i} and $A_{S_{i+1}}$ but in no other set A_{S_j} , $j \in \{1, \dots, l\} - \{i, i+1\}$.

It remains to show that $l \leq 2$. To see this suppose for a contradiction that $l \geq 3$. Define $Z' = \{z, z_1, \dots, z_l\}$ and $\Sigma' = \{S_1, \dots, S_l\}$. Then $\Sigma'_{|Z'}$ is the forbidden split system Γ_{l+1} , a contradiction. \square

Now let S_1 and S_2 be two splits in Σ with the properties given in Lemma 9. Define $C_1 = \{x_1, \dots, x_{b_2}\}$, $D_1 = \{x_{b_2+1}, \dots, x_{b_1-1}\}$, $C_2 = \{x_{b_1}, \dots, x_a\}$ and $D_2 = \{x_{a+1}, \dots, x_n\}$. Select S_1 and S_2 such that $|C_1 \cup C_2|$ is minimal. This induces a bipartition of the split system Σ as described in the following lemma. The routine proof is omitted.

Lemma 10. *Every split in Σ is contained in precisely one of the following subsets of Σ ,*

$$\Sigma_1 = \{S \in \Sigma : C_1 \cup C_2 \cup D_i \subseteq A_S, i \in \{1, 2\}\},$$

$$\Sigma_2 = \{S \in \Sigma : A_S \subseteq C_i \cup D_j, i, j \in \{1, 2\}\}.$$

Next we further study the structure of the splits in Σ_2 . To this end define two elements $p, r \in Z$ to be *clustered*, $p \sim r$, if there exists a split $S \in \Sigma_2$ such that $\{p, r\} \subseteq A_S$. Consider the transitive closure of the binary relation \sim which we denote by the same symbol. The resulting relation \sim is an equivalence relation on Z . Denote the set of equivalence classes with respect to \sim by \mathfrak{F} and call any element in \mathfrak{F} a *cluster*. Note that by construction, for every cluster $F \in \mathfrak{F}$, the split $F|Z - F$ of Z is compatible with ordering Θ . The next lemma concerns the structure of the clusters in \mathfrak{F} .

Lemma 11.

- (a) *For every cluster $F \in \mathfrak{F}$, there exist $i, j \in \{1, 2\}$ such that $F \subseteq C_i \cup D_j$.*
- (b) *There are no two clusters $F_1, F_2 \in \mathfrak{F}$, $F_1 \neq F_2$, such that*
 - (i) $F_1 \cap C_1 \neq \emptyset$, $F_1 \cap D_1 \neq \emptyset$, $F_2 \cap D_1 \neq \emptyset$ and $F_2 \cap C_2 \neq \emptyset$, or
 - (ii) $F_1 \cap C_1 \neq \emptyset$, $F_1 \cap D_2 \neq \emptyset$, $F_2 \cap D_2 \neq \emptyset$ and $F_2 \cap C_2 \neq \emptyset$.

Proof. (a) Assume for contradiction that there exists a cluster $F \in \mathfrak{F}$ that is not contained in $C_i \cup D_j$ for some $i, j \in \{1, 2\}$. The argument can be divided into four very similar cases. We only consider the case that $F \cap D_1 \neq \emptyset$, $F \cap D_2 \neq \emptyset$ and $C_2 \subseteq F$. Then, by the definition of the binary relation \sim , there exist splits $\tilde{S}_1, \dots, \tilde{S}_l$, $l \geq 2$, in Σ_2 and $x_{i_0}, \dots, x_{i_l} \in Z$ such that $x_{i_0} \in D_1$, $\{x_{i_1}, \dots, x_{i_{l-1}}\} \subseteq C_2$, $x_{i_l} \in D_2$, $b_2 + 1 \leq i_0 < i_1 < \dots < i_l \leq n$, and $A_{\tilde{S}_j} \cap \{x_{i_0}, \dots, x_{i_l}\} = \{x_{i_{j-1}}, x_{i_j}\}$ for all $j \in \{1, \dots, l\}$.

Let y be an arbitrary element in C_1 . Then $\{S_1, S_2, \tilde{S}_1, \dots, \tilde{S}_l\}_{|x_{i_0}, \dots, x_{i_l}, z, y}$ is the forbidden split system Γ_{l+3} . Thus, Σ is not clean on X , a contradiction.

(b) We only show (i), then (ii) follows by symmetry. Suppose for contradiction that two clusters $F_1, F_2 \in \mathfrak{F}$, $F_1 \neq F_2$, with property (i) exist. Then, by the definition of the binary relation \sim , there

exist splits \tilde{S}_1, \tilde{S}_2 in Σ_2 and $x_{i_0}, \dots, x_{i_3} \in Z$ such that $x_{i_0} \in C_1$, $\{x_{i_1}, x_{i_2}\} \subseteq D_1$, $x_{i_3} \in C_2$, $1 \leq i_0 < i_1 < i_2 < i_3 \leq a$, $A_{\tilde{S}_1} \cap \{x_{i_0}, \dots, x_{i_3}\} = \{x_{i_0}, x_{i_1}\}$, and $A_{\tilde{S}_2} \cap \{x_{i_0}, \dots, x_{i_3}\} = \{x_{i_2}, x_{i_3}\}$.

Select an arbitrary element $y \in D_2$. Then $\{S_1, S_2, \tilde{S}_1, \tilde{S}_2\}_{\{x_{i_0}, \dots, x_{i_3}, y, z\}}$ is the forbidden split system Ψ , a contradiction. \square

The next lemma helps to simplify the remainder of the proof.

Lemma 12. *Without loss of generality, we can assume that neither $\{x_1, x_n\}$ nor $\{x_{b_1-1}, x_{b_1}\}$ is contained in a cluster in \mathfrak{F} .*

Proof. By Lemma 11(b) at most one of $\{x_1, x_n\}$ and $\{x_a, x_{a+1}\}$ can be contained in a cluster in \mathfrak{F} and, similarly, at most one of $\{x_{b_2}, x_{b_2+1}\}$ and $\{x_{b_1-1}, x_{b_1}\}$ can be contained in a cluster in \mathfrak{F} .

Now consider the case that $\{x_{b_1-1}, x_{b_1}\}$ and $\{x_a, x_{a+1}\}$ are each contained in a cluster in \mathfrak{F} (all other cases can be dealt with similarly). Then we must have that neither $\{x_1, x_n\}$ nor $\{x_{b_2}, x_{b_2+1}\}$ are contained in a cluster in \mathfrak{F} . Furthermore, by Lemma 11(a), there must exist some $c \in \{b_1, \dots, a\}$ such that $\{x_c, x_{c+1}\}$ is not contained in a cluster in \mathfrak{F} . Moreover, by our assumption above, $\{x_{b_2}, x_{b_2+1}\}$ is not contained in a cluster in \mathfrak{F} .

Now it can be checked that every split in $\Sigma|_Z$ is compatible with the ordering

$$\Theta' = x_1, \dots, x_{b_2}, x_c, x_{c-1}, \dots, x_{b_2+1}, x_{c+1}, x_{c+2}, \dots, x_n.$$

So, we could use ordering Θ' instead of ordering Θ and then would have that neither $\{x_1, x_n\}$ nor $\{x_{b_1-1}, x_{b_1}\}$ is contained in a cluster in \mathfrak{F} . \square

Now we construct an ordering of X that is compatible with Σ . This yields a contradiction to the fact that Σ is not circular and finishes the proof. To this end we define

$$Z'_1 = \{x_1, \dots, x_{b_1-1}, y, z\} \quad \text{and} \quad Z'_2 = \{x_{b_1}, \dots, x_n, y, z\}$$

where y is a new element not contained in X . With respect to Z'_1 , the new element y can be thought of as representing an arbitrary element in D_2 . Similarly, with respect to Z'_2 , the new element y can be thought of as representing an arbitrary element in D_1 . Note that $|Z'_1| \leq n$ and $|Z'_2| \leq n$.

Define the bipartitions $\Sigma_1 = \Sigma_1^1 \cup \Sigma_1^2$ and $\Sigma_2 = \Sigma_2^1 \cup \Sigma_2^2$ by

$$\begin{aligned} \Sigma_1^1 &= \{S \in \Sigma_1: D_1 \subseteq A_S\}, & \Sigma_1^2 &= \{S \in \Sigma_1: D_2 \subseteq A_S\}, \\ \Sigma_2^1 &= \{S \in \Sigma_2: A_S \subseteq C_1 \cup D_1\}, & \Sigma_2^2 &= \{S \in \Sigma_2: A_S \subseteq C_2 \cup D_2\}. \end{aligned}$$

For every split $S \in \Sigma$, we define $B_S = X - A_S$. Now we construct a split system Σ'_1 on Z'_1 as follows:

$$\{B_S|Z'_1 - B_S: S \in \Sigma_1^2\} \cup \{A_S|Z'_1 - A_S: S \in \Sigma_1^1\} \cup \{\{y, z\}|Z'_1 - \{y, z\}\}.$$

Similarly, we construct a split system Σ'_2 on Z'_2 :

$$\{B_S|Z'_2 - B_S: S \in \Sigma_1^1\} \cup \{A_S|Z'_2 - A_S: S \in \Sigma_2^2\} \cup \{\{y, z\}|Z'_2 - \{y, z\}\}.$$

Bearing in mind that y can be thought of as an element in D_1 and D_2 , respectively, it follows that the split system Σ'_i is clean on Z'_i , $i \in \{1, 2\}$. Hence, by the minimality of $|X|$, there exists a circular ordering $\Theta'_i = p_1, \dots, p_{l_i}$ of Z'_i that is compatible with Σ'_i . Since the split $\{y, z\}|Z'_1 - \{y, z\}$ is compatible with Θ'_1 we can assume that $p_{l_1-1} = z$ and $p_{l_1} = y$. Similarly, by the minimality of $|X|$, there

exists a circular ordering $\Theta'_2 = r_1, \dots, r_{l_2}$ of Z'_2 that is compatible with Σ'_2 and we can assume that $r_1 = y$ and $r_2 = z$.

Now define the ordering $\tilde{\Theta} = p_1, p_2, \dots, p_{l_1-1}, r_3, r_4, \dots, r_{l_2}$ of X . It is not hard to check that every split in Σ is compatible with $\tilde{\Theta}$. But this contradicts our assumption that Σ is not circular, completing the proof of Theorem 8. \square

Acknowledgments

Author Semple was supported by the New Zealand Marsden Fund. Authors Moulton and Spillner were supported by the Engineering and Physical Sciences Research Council [grant number EP/D068800/1]. Part of this work was completed while the authors participated in the Phylogenetics Programme at the Isaac Newton Institute for Mathematical Sciences. We would also like to thank the anonymous referee for his helpful comments.

References

- [1] H.-J. Bandelt, A. Dress, Reconstructing the shape of a tree from observed dissimilarity data, *Adv. in Appl. Math.* 7 (1986) 309–343.
- [2] H.-J. Bandelt, A. Dress, A canonical split decomposition theory for metrics on a finite set, *Adv. Math.* 92 (1992) 47–105.
- [3] H.-J. Bandelt, A. Dress, Split decomposition: A new and useful approach to phylogenetic analysis of distance data, *Mol. Phylogen. Evol.* 1 (1992) 242–252.
- [4] H.-J. Bandelt, A. Dress, A relational approach to split decomposition, in: O. Opitz, et al. (Eds.), *Information and Classification*, Springer, Berlin, 1993, pp. 123–131.
- [5] D. Bryant, V. Moulton, NeighborNet: An agglomerative method for the construction of phylogenetic networks, *Mol. Biol. Evol.* 21 (2004) 255–265.
- [6] P. Buneman, The recovery of trees from measures of dissimilarity, in: F. Hodson, et al. (Eds.), *Mathematics in the Archaeological and Historical Sciences*, Edinburgh Univ. Press, Edinburgh, 1971, pp. 387–395.
- [7] H. Colonius, H. Schulze, Repräsentation nichtnumerischer Ähnlichkeitsdaten durch Baumstrukturen, *Psych. Beitr.* 21 (1979) 98–111.
- [8] H. Colonius, H. Schulze, Tree structures for proximity data, *British J. Math. Statist. Psych.* 34 (1981) 167–180.
- [9] J. Cracraft, M. Donoghue (Eds.), *Assembling the Tree of Life*, Oxford Univ. Press, Oxford, 2004.
- [10] J. Dopazo, A. Dress, A. von Haeseler, Split decomposition: A technique to analyze viral evolution, *Proc. Natl. Acad. Sci. USA* 90 (1993) 10320–10324.
- [11] A. Dress, P. Erdős, X -trees and weighted quartet systems, *Ann. Combin.* 7 (2003) 155–169.
- [12] J. Felsenstein, *Inferring Phylogenies*, Sinauer, Sunderland, MA, 2003.
- [13] S. Grünewald, K. Forslund, A. Dress, V. Moulton, QNet: An agglomerative method for the construction of phylogenetic networks from weighted quartets, *Mol. Biol. Evol.* 24 (2007) 532–538.
- [14] S. Grünewald, K. Huber, V. Moulton, C. Semple, Encoding phylogenetic trees in terms of weighted quartets, *J. Math. Biol.* 56 (2008) 465–477.
- [15] D. Huson, SplitsTree: Analyzing and visualizing evolutionary data, *Bioinformatics* 14 (1998) 68–73.
- [16] D. Huson, D. Bryant, Application of phylogenetic networks in evolutionary studies, *Mol. Biol. Evol.* 23 (2006) 254–267.
- [17] M. Salemi, A. Vandamme (Eds.), *The Phylogenetic Handbook*, Cambridge Univ. Press, Cambridge, 2003.
- [18] C. Semple, M. Steel, *Phylogenetics*, Oxford Univ. Press, Oxford, 2003.
- [19] K. Strimmer, A. von Haeseler, Quartet puzzling: A quartet maximum likelihood method for reconstructing tree topologies, *Mol. Biol. Evol.* 13 (1996) 964–969.
- [20] R. Wetzel, Zur Visualisierung abstrakter Ähnlichkeitsbeziehungen, PhD thesis, Universität Bielefeld, 1995.
- [21] J. Weyer-Menkoff, New quartet methods in phylogenetic combinatorics, PhD thesis, Universität Bielefeld, 2003.
- [22] J. Weyer-Menkoff, C. Devauchelle, A. Grossmann, S. Grünewald, Integer linear programming as a tool for constructing trees from quartet data, *Comput. Biol. Chem.* 29 (2005) 196–203.